

References

1. Hermentin P, Cuesta-Linker T, Weisse J, Schmidt KH, Knorst M, Scheld M, Thimme M. Comparative analysis of the activity and content of different streptokinase preparations. *Eur Heart J* 2005;26:933–940.
2. Longstaff C, Thelwell C, Whitton C. The poor quality of streptokinase products in use in developing countries. *J Thromb Haemost* 2005; 3: 1092–1093.

Michael Thimme
 ZLB Behring GmbH
 Business Unit Critical Care
 35037 Marburg
 Germany
 Tel: +49 6421 39 4130
 Fax: +49 6421 39 4146
 E-mail address:
 michael.thimme@zlbbehrling.com

doi:10.1093/eurheartj/ehi422

Online publish-ahead-of-print 13 July 2005

Cholesterol levels and cholesterol lowering in idiopathic dilated cardiomyopathy

We read with great interest the recent article of Dr Christ *et al.*¹ about the prognostic significance of serum cholesterol levels in patients with idiopathic dilated cardiomyopathy (IDC). It was found that decreased cholesterol levels do not independently predict adverse prognosis in patients with IDC and that low cholesterol levels are dependent on the severity of the disease (i.e. higher NYHA class, higher left ventricular end-diastolic diameter, and lower left ventricular ejection fraction). In contrast, the authors present an intriguing curve showing that, retrospectively, the prognosis (survival or transplant free survival) was significantly better for the few patients (~15%) who received statins at any time during follow-up. As stated in the discussion, beneficial effects of statins in heart failure patients indeed appear promising in some preliminary results, even in patients without coronary artery disease.² However, the particular analysis and results of Dr Christ may, in our opinion, be related to the fact that the patients who received statins were likely to be those with the highest cholesterol levels, i.e. those with the lower severity of the disease as suggested by the authors themselves. Therefore, it is really not sure that the better prognosis was related to the treatment with statins or to the lipid lowering. Prospective randomized intervention trials are thus particularly needed in the field.

References

1. Christ M, Klima T, Grimm W, Mueller HH, Maisch B. Prognostic significance of serum cholesterol levels in patients with idiopathic dilated cardiomyopathy. *Eur Heart J* 2005. Published online ahead of print April 8, 2005.
2. Krum H, McMurray JJ. Statins and chronic heart failure: do we need a large-scale outcome trial? *J Am Coll Cardiol* 2002;39: 1567–1573.

Laurent Fauchier
 Service de Cardiologie B.
 Centre Hospitalier Universitaire Trousseau
 37044 Tours
 France
 Tel: +33 2 47 47 46 50
 Fax: +33 2 47 47 59 19
 E-mail address: lfau@med.univ-tours.fr

Axel de Labriolle
 Service de Cardiologie A.
 Centre Hospitalier Universitaire Trousseau
 37044 Tours
 France

doi:10.1093/eurheartj/ehi423

Online publish-ahead-of-print 13 July 2005

Cholesterol levels and cholesterol lowering in idiopathic dilated cardiomyopathy: reply

We greatly appreciate the comments of Fauchier *et al.* regarding our recently published findings.¹ Experimental and observational studies have raised concerns whether cholesterol lowering may adversely affect outcome in patients with chronic heart failure. We believe that those studies may not adequately address this clinically important issue due to (i) the artificial nature of those experimental studies and (ii) the mixed patient populations analysed including patients with coronary artery disease, which is an inflammatory disease by itself.

Owing to the exceptional opportunity in our institution, where patients with idiopathic dilated cardiomyopathy (iDCM) have been followed for several years (Marburg cardiomyopathy database), we have been able to examine aforementioned questions in a large cohort of iDCM patients, in whom systolic dysfunction due to other causes has been included on thorough examinations. Our data of this well-characterized patient group convincingly demonstrate that cholesterol levels depend on the severity of cardiac disease. Surprisingly, further analysis

indicates that statins may beneficially affect outcome in iDCM patients supporting experimental and preliminary clinical findings in this field.^{2,3} Although our data appear conclusive, they were derived from *post hoc* analysis with all limitations inherent with subgroup analysis. In addition, (i) a large number of patients on statins did not receive those drugs throughout the whole observational period; (ii) it cannot be excluded that iDCM patients with high cholesterol levels, which is associated with better outcome as already demonstrated in our study, are preferentially treated with statins leading to bias, although we have tried to adjust during statistical analysis. Subsequently, our data need to be confirmed by well-designed, prospective, randomized trials as suggested by Fauchier. Randomized clinical studies are already on the way to clarify this exciting concept (<http://cvm.controlled-trials.com/content/2/6/266>).

References

1. Christ M, Klima T, Grimm W, Mueller HH, Maisch B. Prognostic significance of serum cholesterol levels in patients with idiopathic dilated cardiomyopathy. *Eur Heart J* 2005. Published online ahead of print April 8, 2005.
2. Bauersachs J, Galuppo P, Fraccarollo D, Christ M, Ertl G. Improvement of left ventricular remodeling and function by hydroxymethylglutaryl coenzyme A reductase inhibition with cerivastatin in rats with heart failure after myocardial infarction. *Circulation* 2001;104: 982–985.
3. Laufs U, Wassmann S, Schackmann S, Heesch C, Bohm M, Nickenig G. Beneficial effects of statins in patients with non-ischemic heart failure. *Z Kardiol* 2004;93:103–108.

Michael Christ
 Department of Internal Medicine
 Medical Division A
 University Hospital Basel
 University of Basel
 Petersgraben 4
 CH-4031 Basel
 Switzerland
 Tel: +41 61 265 2525
 Fax: +49 69 13303154027
 E-mail address: Micchrist@uhbs.ch

and
 Department of Internal Medicine, Cardiology
 Philipps University Marburg
 Germany

Theresia Klima
 Department of Internal Medicine
 Medical Division A
 University Hospital Basel
 University of Basel
 Petersgraben 4
 CH-4031 Basel
 Switzerland
 and

Department of Internal Medicine,
Cardiology
Philipps University Marburg
Germany

Wolfram Grimm
Department of Internal Medicine,
Cardiology
Philipps University Marburg
Germany

doi:10.1093/eurheartj/ehi434

Online publish-ahead-of-print 25 July 2005

Combined pharmacologic treatment with clopidogrel and statin for patients with acute coronary syndrome: is there a survival advantage?

We read with great interest the retrospective study by Lim *et al.*¹ on the impact of combined treatment of anti-platelet drugs and statin on the clinical outcomes of patients with acute coronary syndrome (ACS). The authors in their study suggest that despite the possibility of competitive drug interaction between statin and clopidogrel,² the combination of clopidogrel and statin has beneficial synergetic effects on the mortality of patients with non-ST-segment elevation ACS.

Randomized controlled trials over the past 20 years clearly show that aspirin, beta-blocker, ACE-inhibitor, and statin therapies prevent recurrent coronary events in patients with ACS.^{3,4} In particular, statins have been associated with a substantial reduction of vascular morbidity and mortality. The benefits of statin therapy have been demonstrated in patients with coronary artery disease, hypertension, diabetes, and in those with elevated, averaged,⁵ or even near-normal LDL-cholesterol levels.⁶ Statins are known to promote an overall 24–37% reduction in the relative risk for coronary events, independent of other confounding variables such as age and sex (for review refer to Gotto⁷). Thus, statins are recommended in the current therapeutic guidelines for the treatment of ACS.^{3,4}

Data provided by Lim *et al.*¹ show that the 6-month mortality estimated by the Kaplan-Meier curve was significantly lower in group IV (aspirin + clopidogrel + statin) compared with group II (aspirin + clopidogrel). We would like to make the following important notes: (1) the benefits of statins, beta-blockers, and ACE-inhibitors on the

morbidity and mortality of patients with ACS have been well documented, (2) a statistically significant greater number of group IV patients compared with group II were on beta-blockers and ACE-inhibitors in the study conducted by Lim *et al.*,¹ and (3) *ex vivo* studies of clopidogrel and statin drug interactions show significant inhibition of clopidogrel metabolism but not statin efficacy.

Therefore, this suggests that the low mortality observed by Lim *et al.*¹ within group IV compared with group II could be simply attributed to the independent statin effect and/or to the effect of beta-blockers and ACE-inhibitors. The Lim *et al.*¹ study shows that statin significantly reduces the mortality in the 'treated' group IV compared with the 'non-treated' group II, and aspirin + clopidogrel could be considered as a background for both groups. In addition, the mortality rate of group III (aspirin + statin) and group II (aspirin + clopidogrel) were 3.6 and 4.4%, respectively. This further indicates the value of statin therapy in patients with ACS.

The Kaplan-Meier curve in the Lim *et al.*¹ study shows only the unadjusted mortality for group II vs. group IV. 'Restricting the main study comparisons to the two groups in whom clopidogrel was used', as stated by the authors, might not be sufficient to evaluate the synergistic effect of the drugs of interest on overall mortality. To investigate the additive and/or synergetic effect of clopidogrel and statin, Kaplan-Meier plot should include the adjusted mortality of all the four groups studied by Lim *et al.*

In conclusion, the retrospective study conducted by Lim *et al.*¹ in ACS patients makes it difficult to draw the conclusion that a 'clinically' superior effect exists for clopidogrel-statin drug combination. Moreover, the conclusion of additive or synergetic effects of the combined two drugs on the clinical outcomes of patients with ACS warrants further investigation.

References

1. Lim MJ, Spencer FA, Gore JM, Dabbous OH, Agnelli G, Kline-Rogers EM, Dibenedetto D, Eagle KA, Mehta RH. Impact of combined pharmacologic treatment with clopidogrel and a statin on outcomes of patients with non-ST-segment elevation acute coronary syndromes: perspectives from a large multinational registry. *Eur Heart J* 2005;26: 1063–1069.
2. Lau WC, Waskell LA, Watkins PB, Neer CJ, Horowitz K, Hopp AS, Tait AR, Carville DGM, Guyer KE, Bates ER. Atorvastatin reduces the ability of clopidogrel to inhibit platelet aggregation: a new drug-drug interaction. *Circulation* 2003;107:32–37.
3. Antman EM, Anbe DT, Armstrong PW, Bates ER, Green LA, Hand M, Hochman JS, Krumholz HM, Kushner FG, Lamas GA, Mullany CJ, Ornato JP, Pearle DL, Sloan MA, Smith SC Jr, Alpert JS, Anderson JL, Faxon DP, Fuster V, Gibbons RJ, Gregoratos G, Halperin JL, Hiratzka LF, Hunt SA, Jacobs AK; American College of Cardiology; American Heart Association Task Force on Practice Guidelines; Canadian Cardiovascular Society. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1999 Guidelines for the Management of Patients with Acute Myocardial Infarction). *Circulation* 2004;110:e82–e292.
4. Bertrand ME, Simoons ML, Fox KA, Wallentin LC, Hamm CW, McFadden E, De Feyter PJ, Specchia G, Ruzylo W; Task Force on the Management of Acute Coronary Syndromes of the European Society of Cardiology. Management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 2002;23:1809–1840.
5. Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, Cole TG, Brown L, Warnica JW, Arnold JM, Wun CC, Davis BR, Braunwald E. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med* 1996; 335:1001–1009.
6. The HPS Study Group. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals. *Lancet* 2002;360:7–22.
7. Gotto AM. Statin therapy: where are we? Where do we go next? *Am J Cardiol* 2001;87(suppl.): 13B–18B.

Zakaria Almsheerqi

Department of Physiology
Faculty of Medicine
National University of Singapore
Block MD9

2 Medical Drive
Singapore 117597

Tel: +65 9237 7529

Fax: +65 6778 8161

E-mail address: phszama@nus.edu.sg

Craig McLachlan and Peter J. Mossop

Department of Medical Imaging
St Vincents Hospital
Melbourne
Australia

Yuru Deng

Department of Physiology
Faculty of Medicine
National University of Singapore
Block MD9
2 Medical Drive
Singapore 117597
Singapore